



This document is sponsored by  
**The Science Foundation College** Kiwanga- Namanve  
 Uganda East Africa  
 Senior one to senior six  
 +256 778 633 682, 753 802709  
**Based on, best for sciences**




## Vectors

A vector is a quantity which has both magnitude and direction. Examples include, force, displacement, acceleration, momentum and velocity.

### Representation of a vector


A vector is represented by a line with an arrow to indicate the direction of the vector.



where the order of the letters shows the direction

### Vectors in dimensions

Vectors can be represented in three dimensions as i, j and k along the x, y and z- axes respectively



### Resultant of vectors

When several vectors ( $V_1, V_2, V_3... V_n$ ) are acting on a point object, the net vector R, is calculated as the vector sum

$$R = V_1 + V_2 + V_3 + \dots + V_n = \sum_{r=1}^n V_r$$

#### Example 1

Find the resultant of the following vectors

- (a)  $(2i + 3j + 3k)$  and  $(2i + 4j - 8k)$

$$R = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} + \begin{pmatrix} 2 \\ 4 \\ -8 \end{pmatrix} = \begin{pmatrix} 4 \\ 7 \\ -5 \end{pmatrix} \text{ or } 4i + 7j - 5k$$

- (b)  $(7i - 4j + 3k)$ ,  $(5i - 2j + 8k)$ ,  $(i - k)$

$$R = \begin{pmatrix} 7 \\ -4 \\ 3 \end{pmatrix} + \begin{pmatrix} 5 \\ -2 \\ 8 \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} 13 \\ -6 \\ 10 \end{pmatrix} \text{ or } 13i - 6j + 10k$$

#### Example 2

The resultant of  $(5i - 2j)$ ,  $(7i + 4j)$ ,  $(ai + bj)$  and  $(-3i + 2j)$  is  $(5i + 5j)$ . Find the values of a and b.

$$R = \begin{pmatrix} 5 \\ -2 \end{pmatrix} + \begin{pmatrix} 7 \\ 4 \end{pmatrix} + \begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} -3 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}$$

$$= \begin{pmatrix} 9 + a \\ 4 + b \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}$$

$$9 + a = 5; a = -4 \text{ also } 4 + b = 5; b = 1$$

### Example 3

The resultant of the forces  $(3i + (a-c)j)N$ ,  $((2a + 3c)i + 5j)N$  and  $(4i, 6j)N$  acting on a particle is  $(10i + 12j)N$ . find

(i) Values of a and c

$$R = \begin{pmatrix} 3 \\ a - c \end{pmatrix} + \begin{pmatrix} 2a + 3c \\ 5 \end{pmatrix} + \begin{pmatrix} 4 \\ 6 \end{pmatrix} = \begin{pmatrix} 10 \\ 12 \end{pmatrix}$$

$$2a + 3c + 7 = 10$$

$$2a + 3c = 3 \dots\dots\dots (i)$$

$$a - c + 11 = 12$$

$$a - c = 1 \dots\dots\dots (ii)$$

$$(i) + 3(ii)$$

$$5a = 6; a = 1.2$$

from eqn. (ii)

$$c = 0.2$$

(ii) magnitude of  $(2a + 3c)i + 5j$

$$R = (2a + 3c)i + 5j = (1.2 \times 2 + 3 \times 0.2)i + 5j = 3i + 5j$$

$$|R| = \sqrt{3^2 + 5^2} = 5.831N$$

### Magnitude or modulus of a vector

This is the length of a vector

(i) Given  $R = xi + yj$ ;  $|R| = \sqrt{x^2 + y^2}$

(ii) Given  $R = xi + yj + zk$ ;  $|R| = \sqrt{x^2 + y^2 + z^2}$

### Example 5


Find the magnitude of the following vectors

(a)  $3i + 4j$ ;  $|R| = \sqrt{3^2 + 4^2} = 5$

(b)  $3i + 2j - 6k$ ;  $|R| = \sqrt{3^2 + 2^2 + (-6)^2} = 7$

### Direction of a vector

Consider  $R = xi + yj$



$$\theta = \tan^{-1} \left( \frac{y}{x} \right)$$


### Example 6

Find the magnitude and direction of the resultant of each of the following

$$(a) (2i + 3j)N, (5i - 2j)N, (-3i, 3j)$$

$$R = \begin{pmatrix} 2 \\ 3 \end{pmatrix} + \begin{pmatrix} 5 \\ -2 \end{pmatrix} + \begin{pmatrix} -3 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$|R| = \sqrt{4^2 + 4^2} = 5.6569N$$




$$\theta = \tan^{-1}\left(\frac{4}{4}\right) = 45^\circ$$

$$(b) \begin{pmatrix} 2 \\ 4 \end{pmatrix} N, \begin{pmatrix} -6 \\ -5 \end{pmatrix} N \text{ and } \begin{pmatrix} 2 \\ 1 \end{pmatrix} N$$

$$R = \begin{pmatrix} 2 \\ 4 \end{pmatrix} + \begin{pmatrix} -6 \\ -5 \end{pmatrix} + \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \end{pmatrix}$$

$$|R| = \sqrt{(-2)^2 + 0^2} = 2N$$




$$\theta = \tan^{-1}\left(\frac{0}{-2}\right) = 180^\circ$$

### Example 7

Four forces of  $ai + (a-1)j$ ,  $3i + 2aj$ ,  $5i - 6j$ , and  $-i - 2j$  act on a particle. The resultant forces make an angle of  $45^\circ$  with horizontal. Find  $a$ . Hence determine the magnitude of the resultant force.

$$R = \begin{pmatrix} a \\ a-1 \end{pmatrix} + \begin{pmatrix} 3 \\ 2a \end{pmatrix} + \begin{pmatrix} 5 \\ -6 \end{pmatrix} + \begin{pmatrix} -1 \\ -2 \end{pmatrix}$$

$$= \begin{pmatrix} a+7 \\ 3a-9 \end{pmatrix}$$



$$\frac{a+7}{3a-9} = \tan^{-1}(45^\circ) = 1$$

$$a+7 = 3a-9, a = 8$$

$$R = \begin{pmatrix} a+7 \\ 3a-9 \end{pmatrix} = \begin{pmatrix} 8+7 \\ 3 \times 8 - 9 \end{pmatrix} = \begin{pmatrix} 15 \\ 15 \end{pmatrix}$$

$$|R| = \sqrt{15^2 + 15^2} = 21.21N$$

### Unit vector

This a vector whose magnitude is unit (1)

$$\text{Unit vector of } r \text{ denoted by } \hat{r} = \frac{r}{|r|}$$

### Example 8

Find the unit vector of  $a = 6i - 2j + 3k$

Solution

$$\hat{a} = \frac{6i - 2j + 3k}{\sqrt{6^2 + (-2)^2 + 3^2}} = \frac{1}{7}(6i - 2j + 3k)$$

Parallel vectors

If vectors  $a$  and  $b$  are parallel, then one of them is a scalar multiple of the other.

$$\text{If a vector } r \text{ of magnitude } |r| \text{ moves in direction } xi + yj + zk \text{ then, } r = |r| \left( \frac{xi + yj + zk}{\sqrt{x^2 + y^2 + z^2}} \right)$$

### Example 9

Find the vector,  $V$  which has a magnitude of 15 units and is parallel to  $16i + 12j$

$$V = 15 \times \frac{16i + 12j}{\sqrt{16^2 + 12^2}} = 15 \times \frac{16i + 12j}{20} = 12i + 9j$$

### Example 10

A body of velocity  $v$  and of magnitude 20m/s moves in the direction  $6i + 8j$ . Find  $V$ .

$$V = 20 \times \frac{6i + 8j}{\sqrt{6^2 + 8^2}} = 20 \times \frac{6i + 8j}{10} = 12i + 16j$$

Example 11

A force of magnitude 12N acts on a body in the direction  $2i + j + 2k$ . Find the force

$$V = 12 \times \frac{2i + j + 2k}{\sqrt{2^2 + 1^2 + 2^2}} = 12 \times \frac{2i + j + 2k}{3} = 8i + 4j + 8k$$

**Example 12**

The force A of magnitude 5N in the direction with unit vector  $\frac{1}{5}(3i + 4j)$  and force B of magnitude 13N in the direction with unit vector  $\frac{1}{13}(5i - 12j)$ . Find the resultant forces of A and B.

$$\begin{array}{l} F_A = \frac{1}{5}(3i + 4j) \times 5 = 3i + 4j \\ F_B = \frac{1}{13}(5i - 12j) \times 13 = 5i - 12j \end{array} \quad \left| \begin{array}{l} F = \begin{pmatrix} 3 \\ 4 \end{pmatrix} + \begin{pmatrix} 5 \\ -12 \end{pmatrix} = \begin{pmatrix} 8 \\ -8 \end{pmatrix} \\ |F| = \sqrt{8^2 + (-8)^2} = 11.3137\text{N} \end{array} \right.$$

**Example 13**

A particle P moves through a displacement of 2m when acted on by two forces  $F_1$  and  $F_2$ . Find the work done by the resultant force, if  $F_1 = i - j$  and  $F_2 = 10 \text{ N}$  and acts in the direction  $4i + 3j$

Solution

$$F_1 = i - j$$

$$F_2 = 10 \times \frac{4i + 3j}{\sqrt{4^2 + 3^2}} = 8i + 6j$$

$$\left| \begin{array}{l} F = \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \begin{pmatrix} 8 \\ 6 \end{pmatrix} = \begin{pmatrix} 9 \\ 5 \end{pmatrix} \\ |F| = \sqrt{9^2 + 5^2} = 10.2956\text{N} \\ W = |F| \times d = 10.2956 \times 2 = 20.5912\text{J} \end{array} \right.$$

### Revision exercise 1

- Find the resultant of each of the following forces
  - $(6i + 2j)\text{N}$ ,  $(-5i + j)\text{N}$ ,  $(3i - 3j)\text{N}$ .  $[(4i)\text{N}]$
  - $(2i + 4j)\text{N}$ ,  $(3i - 5j)$ ,  $(6i + 2j)\text{N}$ ,  $(-7i - 7j)\text{N}$ .  $[(4i - 6j)\text{N}]$
  - $(2i + 3j - 7k)\text{N}$ ,  $(2i + 5k)\text{N}$ ,  $(3j + 4k)\text{N}$ .  $[(4i + 6j + 2k)\text{N}]$
- The resultant of forces  $(5i + 7j)$ ,  $(ai + bj)$  and  $(bi - aj)\text{N}$  is a force  $(11i + 5j)\text{N}$ . Find a and b.  
 $[a = 4, b = 2]$
- Find the magnitude and direction of the resultant of each of the following;
  - $(-2i + 5j)\text{N}$ ,  $(i + 2j)\text{N}$ .  $[7.07\text{N at } 98.1^\circ]$
  - $(6i + 2j)\text{N}$ ,  $(4i - 3j)\text{N}$ .  $[10.05\text{N at } 354.3^\circ]$
  - $(3i + 2j)$ ,  $(-5i + j)\text{N}$ .  $[3.61\text{N at } 124^\circ]$
- A force of magnitude 50N acts on a body in the direction  $24i + 7j$ . Find the force.  $[(48i + 14j)]$
- Two forces  $F_1$  and  $F_2$  have magnitude  $\alpha\text{N}$  and  $\beta\text{N}$  and act in the direction  $i - 2j$  and  $4i + 3j$  respectively. Given that the resultant of  $F_1$  and  $F_2$  is  $(48i + 14j)$ . Find the magnitude of  $\alpha\text{N}$  and  $\beta\text{N}$ .  $[\alpha = 8\sqrt{5}\text{N and } \beta = 50\text{N}]$
- If  $a = 3i + 4j$ ,  $b = 4i + 20j$  and  $c = 5i - 19j$ ; find the
  - resultant of a and b  $[(7i + 24j)]$
  - resultant of a and c  $[(8i - 15j)]$
  - vector is parallel to a and has magnitude of 15 unit  $[(9i + 12j)]$

- (iv) vector parallel to  $(a + b)$  and has a magnitude of 100 units  $[(28i + 96j)]$
7. If  $a = 2i + 5j$ ,  $b = -7i + 7j$  and  $14i$ . Find the;
- resultant of  $a$  and  $b$   $[(-5i + 12j)]$
  - resultant of  $a$ ,  $b$  and  $c$   $[(9i + 12j)]$
  - $|b|$   $[7\sqrt{2}]$
  - $|a + b + c|$   $[15\text{units}]$
  - vector is parallel to  $a$  and has a magnitude of  $5\sqrt{29}$  units.  $(10i + 25j)$
  - Vector is parallel to  $(a + b + c)$  and has magnitude 90 units.  $[(54i + 72j)]$
8. If  $a = i - 3j + 2k$ ,  $b = 5i + 4j$  and  $c = 3i + j + 4k$ . Find the
- resultant of  $a$  and  $b$   $[(6i + j + 2k)]$
  - resultant of  $a$ ,  $b$  and  $c$ .  $[(9i + 2j + 6k)]$
  - $|a|$   $[\sqrt{14}]$
  - $|a + b + c|$   $[11\text{units}]$
  - Vector parallel to  $(a + b + c)$  and has magnitude 5 units  $[\frac{5}{11}(9i + 2j + 6k)]$
9. If  $a = 2i + 7j + 7k$ ,  $b = 6i - 3j + 2k$  and  $c = -4j - 3k$ . find the
- resultant  $a$  and  $b$   $[8i + 4j + 9k]$
  - resultant  $a$  and  $c$   $[2i + 3j + 4k]$
  - $|b|$   $[7\text{units}]$
  - $|a + b + c|$   $[10\text{ units}]$
  - vector is parallel to  $|a + b + c|$  and has magnitude of 50 units  $[40i + 30k]$

### Scalar products or dot products

The dot product of two vectors  $a$  and  $b$  inclined at angle  $\theta$  is given by

$$a \cdot b = |a||b| \cos\theta$$

#### Note

If two vectors are perpendicular then the angle between them is  $90^\circ$  and

$$a \cdot b = |a||b| \cos 90 = 0$$

#### Example 14

If  $a = i - 2k$  and  $b = 3i - 3j + k$ , find

- $a \cdot b$
- the angle between  $a$  and  $b$

Solution

$$(i) \quad a \cdot b = \begin{pmatrix} 1 \\ 0 \\ -2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -3 \\ 1 \end{pmatrix} = 3 + 0 + -2 = 1$$

$$(ii) \quad \theta = \cos^{-1} \left( \frac{a \cdot b}{|a||b|} \right) = \cos^{-1} \frac{1}{\sqrt{1^2 + (2)^2} \sqrt{3^2 + (-3)^2 + 1^2}} = 84.1^\circ$$

#### Example 15

If  $p = 2i - j + 3k$  and  $q = i + 4j + 3k$ ; find the angle between  $p$  and  $q$ .

Solution

$$p \cdot q = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 4 \\ 3 \end{pmatrix} = 2 + -4 + 9 = 7$$

$$\theta = \cos^{-1} \left( \frac{p \cdot q}{|p||q|} \right) = \cos^{-1} \frac{7}{\sqrt{2^2 + (-1)^2 + 3^2} \sqrt{1^2 + 4^2 + 3^2}} = 68^\circ$$

### Example 16

If the angle between two vectors  $a = xi + 2j$  and  $b = 3i + j$  is  $45^\circ$ . Find the two possible values of constant  $x$ .

Solution

$$\begin{aligned} \begin{pmatrix} x \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 1 \end{pmatrix} &= \sqrt{x^2 + 2^2} \cdot \sqrt{3^2 + 1^2} \\ 3x + 2 &= \sqrt{x^2 + 2^2} \cdot \sqrt{10} \cdot \frac{\sqrt{2}}{2} \\ (3x + 2)^2 &= (x^2 + 4) \cdot 10 \cdot \frac{2}{4} \end{aligned} \quad \left| \begin{array}{l} x^2 + 3x - 4 = 0 \\ (x + 4)(x - 1) = 0 \\ x = -4 \text{ and } x = 1 \end{array} \right.$$

### Example 17

If  $p = 2ai + 7j - k$  and  $q = 3ai + \alpha j + 3k$ . Find the value of the scalar  $\alpha$  if the vectors are perpendicular

Solution

$$\begin{aligned} \begin{pmatrix} 2\alpha \\ 7 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 3\alpha \\ \alpha \\ 3 \end{pmatrix} &= 0 \\ 6\alpha^2 + 7\alpha - 3 &= 0 \\ \alpha &= \frac{1}{3} \text{ and } \alpha = \frac{3}{2} \end{aligned}$$

## Revision exercise 2

1. Find the scalar products for each of the following pairs of vectors.

(i) $a = 2i + j, b = i - 3j$ [-1]	$\begin{aligned} &\begin{pmatrix} 0 \\ 5 \\ -2 \end{pmatrix} \text{ and } \begin{pmatrix} -3 \\ 2 \\ 1 \end{pmatrix} [8] \\ &\begin{pmatrix} 5 \\ 2 \\ 7 \end{pmatrix} \text{ and } \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} [2] \end{aligned}$
(ii) $a = 3i, b = -2i + j$ [-6]	
(iii) $a = 5i + j - 2k, b = 4i + 3j - 8k$ [39]	
(iv) $2i + 4j - 15k$ and $-8i + 2j - k$ [7]	
(v) $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 3 \\ -2 \end{pmatrix}$ [4]	
(vi) $\begin{pmatrix} 5 \\ -1 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 4 \end{pmatrix}$ [6]	

2. Find the angles between each of the following pairs of vectors

(i) $3i + 4j$ and $5i - 12j$ [ $121^\circ$ ]	$\begin{aligned} &\begin{pmatrix} 3 \\ 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 \\ -2 \end{pmatrix} [82^\circ] \\ &\begin{pmatrix} 6 \\ -8 \end{pmatrix} \text{ and } \begin{pmatrix} 5 \\ 4 \end{pmatrix} [92^\circ] \end{aligned}$
(ii) $3i$ and $-2j$ [ $90^\circ$ ]	
(iii) $2i + 3j - 6k$ and $2i + j + 2k$ [ $10^\circ$ ]	
(iv) $i + 2j - k$ and $-i + 2j - k$ [ $48^\circ$ ]	

(v) (vii)  $\begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$  and  $\begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$  [120°]      (viii)  $\begin{pmatrix} 2 \\ 2 \\ 3 \end{pmatrix}$  and  $\begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$  [73°]

3. If  $a = \alpha i + 2j - k$  and  $b = 5i - \alpha j + k$ . Find the value of the scalar  $\alpha$  if the vectors are perpendicular  $[\frac{1}{3}]$
4. If  $a = 2i + \alpha j$  and  $b = -\alpha - k$ . Find the value of the scalar  $\alpha$  if the vectors are perpendicular [0]
5. If  $a = 4i + 5j$  and  $b = qi - 8j$ . Find the value of scalar  $q$  if the vectors are perpendicular. [10]
6. If  $a = 6i - j$  and  $b = 2i + pk$ . Find the value of scalar  $p$  if the vectors are perpendicular [12]
7. Given  $\begin{pmatrix} q \\ 2 + q \\ 3 \end{pmatrix}$  and  $\begin{pmatrix} -1 \\ 3 \\ 4 - q \end{pmatrix}$  are perpendicular vectors. Find the value of  $q$ . [18]
8. If  $a = qi + 8j + (3q + 1)k$  and  $b = (q+1)i + (q-1)j - 2k$ . Find the value of the possible values of constant  $q$  if the vectors are perpendicular. [2 or -2]

Thank you

Dr. Bbosa Science